



Potential environment effect on ultrahigh resolution Sr/Ca of giant clam shells from South China Sea

Chengcheng Liu^{1,2} · Hong Yan^{1,3} · Liqiang Zhao⁴ · Nanyu Zhao¹ · Fan Luo² · Hanfeng Wen¹ · Haotian Yang³ · Wei Yang⁵ · Jialong Hao⁵ · Chengde Liang¹ · Kentaro Tanaka⁶ · Naoko Murakami-Sugihara⁶ · Kotaro Shirai⁶ · Naoto Takahata⁶ · John Dodson^{1,7} · Bernd R. Schöne⁸

Received: 23 February 2024 / Accepted: 21 August 2024 / Published online: 2 September 2024
© The Author(s), under exclusive licence to International Coral Reef Society (ICRS) 2024

Abstract The hourly to monthly resolution Sr/Ca profiles of modern juvenile giant clam (*Tridacna* spp.) shells from the northern South China Sea were obtained using ICP-OES, LA-ICP-MS and NanoSIMS. The results showed the variability of Sr/Ca profiles determined by different analytical methods were consistent on monthly time scale. The hourly resolved Sr/Ca determined by NanoSIMS showed pronounced daily cycles which may be associated with the diurnal cycle of the physiology, environment and/or weather condition. By comparison of daily resolution Sr/Ca and contemporaneous instrumental environment data, no robust link was observed between environmental parameters and juvenile *Tridacna* Sr/Ca, and only weak correlation was found between daily outgoing longwave radiation

(OLR) ($r = -0.167$, $p < 0.01$) and Sr/Ca. However, the effective solar radiation cannot explain the variability of nightly Sr/Ca, and some other factors are more likely to influence the Sr/Ca of juvenile *Tridacna*. Moreover, no consistent correlation was observed between Sr/Ca and growth rate of *Tridacna*. The mechanism of juvenile *Tridacna* Sr/Ca is still mysterious, conducting artificial culture experiments seems the best way to unravel the mechanisms behind Sr²⁺ incorporation into *Tridacna* shells.

Keywords Giant clam shell · Sr/Ca · South China sea · NanoSIMS

Introduction

Precisely dated, ultra-high-resolution (hourly, daily) climate archives could greatly improve our understanding of past environmental changes and enable paleoweather analysis (Saenger et al. 2009; Sano et al. 2012; Cobb et al. 2013; McGregor et al. 2013; Yan et al. 2015, 2020). While trees can provide annually resolved climatic data of the terrestrial realm (Briffa et al. 1990; Esper et al. 2002; Masson-Delmotte et al. 2005; Miller et al. 2006; Fowler et al. 2012, St. George 2014), shallow-water corals and many marine bivalves serve as subseasonally to annually resolved paleoenvironmental recorders in the marine realm (Schöne et al. 2002; Saenger et al. 2009; Cobb et al. 2013; Loubere et al. 2013; McGregor et al. 2013; Dee et al. 2020; García-Escárcaga et al. 2020; Nicastro et al. 2020). However, few paleoclimate archives provide information that would be required for the reconstruction of past weather phenomena, which is still almost blank. One of such archives is the shell of giant clams, e.g., *Tridacna* spp.

✉ Hong Yan
yanhong@ieecas.cn

- 1 State Key Laboratory of Loess and Quaternary Geology, Institute of Earth Environment, Chinese Academy of Sciences, Xi'an 710061, China
- 2 Xi'an Institute for Innovative Earth Environment Research, Xi'an 710061, China
- 3 Institute of Global Environmental Change, Xi'an Jiaotong University, Xi'an 710049, China
- 4 College of Fishery, Guangdong Ocean University, Zhanjiang 524088, China
- 5 Key Laboratory of Earth and Planetary Physics, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing 100029, China
- 6 Atmosphere and Ocean Research Institute, University of Tokyo, Chiba, Japan
- 7 School of Earth, Atmospheric and Life Sciences, University of Wollongong, Wollongong, NSW 2500, Australia
- 8 Institute of Geosciences, University of Mainz, Mainz, Germany